



Assessing Empathy Levels and Their Associations Among 3rd-Year Medical Students at Khyber Medical College, Pakistan

Mashal Khan¹, Hooria Asad¹, Naeemullah², Muhammad Imran Marwat³

¹Khyber Medical College, Peshawar Pakistan

²Department of Community Medicine, Saidu Medical College, Swat

³Department of Community Medicine and Public Health, Khyber Medical College, Peshawar, Pakistan

ABSTRACT

OBJECTIVE: Empathy is a critical factor in doctor-patient relationship. There is well-documented evidence that empathy falls alarmingly during 3rd-year of medical school. Our aim is, therefore, to assess their empathy levels and their associations.

METHODOLOGY: A cross-sectional study was conducted from December 2022 to May 2023 using Jefferson Scale of Empathy student-version (JSE-S) at Khyber Medical College, Pakistan to assess 3rd-year students' empathy. A sample size of 152 was calculated. Data was collected using convenience sampling technique. Descriptive statistics were calculated for JSE-S scores with comparisons across variables using an independent sample t-test and one-way analysis of variance (ANOVA) with post hoc test.

RESULTS: After a response rate of 84.9%, 123 forms (65 males, 58 females) were analyzed. Majority (73.2%) had age < 22 years. Overall, mean empathy score recorded was 103.24 ± 15.83 (range = 53-132). Empathy was significantly higher in group younger than 22 years ($p = 0.004$) and females ($p < 0.001$). The difference was non-significant among specialty preference groups ($p = 0.08$) but separately for females, it was significant ($p = 0.02$), with the underlying difference between technology-oriented specialties and undecided group ($p = 0.02$) on post-hoc comparisons.

CONCLUSION: This article looks into the empathy of 3rd-year medical students. Empathy, as indicated, varied significantly among age and gender groups while specialty preference was found to have little effect. Interestingly, empathy was significantly different for specialty preference only among females. Integration of empathy programs into curriculum is recommended.

KEY WORDS: Empathy, Emotions, Medical student, Patient Care, Pakistan.

*For Correspondence

Dr. Muhammad Imran Marwat

Assistant Professor, Department of Community Medicine and Public Health, Khyber Medical College, Peshawar, Pakistan

Email: imranmarwat_30@kmc.edu.pk

Submission date: 19th June 2023

Acceptance date: 31st October 2023

Publication date: 31st December 2023

INTRODUCTION

Empathy in a clinical setup has four fundamental aspects which has been understood over the years. i.e., Emotive (the capacity to imagine the patient's emotional state), Moral (internally motivated to empathize), Cognitive (the capability to recognize and comprehend patient's emotions), and Behavioral (the capacity to communicate this comprehension of their emotions back to them)¹.

Medical school provides a good platform to acquire empathic behavior which later on is critical in the doctor-patient relationship. Although a moderate level of it towards the patient is necessary for doctors, a certain degree of detachment is also important. Over-attachment and over-detachment both can

have serious consequences for the doctor as well as the patient². Thus, doctors need to be emotionally self-aware to maintain a balanced amount of attachment with their patients.

Many scoring systems of empathy are available like the Interpersonal reactivity index, Toronto empathy questionnaire, Balanced emotional empathy test, Kiersma-Chen empathy scale, and Jefferson scale of physician empathy. Considering the suitability and relatedness of these different instruments, the medical student version of Jefferson scale of empathy has been chosen for this study.

In the literature available, empathy has been correlated with many factors like gender^{3, 4}, year of study³, academic performance⁴, emotional intelligence, gratitude⁵, change along the medical course^{6,7}, and clinical outcomes⁸. But results of these

This Article may be Cited as. Khan M, Asad H, Ullah N, Marwat IM. Assessing Empathy Levels and their Associations Among 3rd-Year Medical Students at Khyber Medical College, Pakistan. Adv Basic Med Sci.2023;7(2)68-73.DOI: <https://doi.org/10.35845/abms.2023.2.286>

studies are highly inconsistent. However, an alarming decline is usually seen in empathy during the 3rd-year of medical school, for which it was termed as the devil's year of medical school⁹, even though this is the period where it is most essential since they have their first clinical exposure.

Empathy among medical students has been assessed around the globe including many parts of Pakistan. But no study of this kind has been carried out in Khyber Medical College. This study intends to generate data regarding this topic to fill the gap in present information.

METHODOLOGY

A cross-sectional study was carried out among 3rd-year students of Khyber Medical College from December 2022 to May 2023. The study was approved by the Institutional Research and Ethical Review Board (IREB) of Khyber Medical College, Peshawar Pakistan under reference no. 888/DME/KMC. All participants gave written informed consent prior to the study. Strobe guidelines were followed here.

Participants and sampling technique: Sample size was computed with the help of online calculator of Raosoft Inc. (http://www.raosoft.com/sample_size.html). Keeping the margin of error and confidence interval at 5% and 95% respectively, and a population size of 250 with a response distribution of 50% which is a conservative assumption, the recommended sample size was obtained to be 152. In Pakistan, MBBS is a 5-year study plan having the first two years devised as pre-clinical years, which are followed by three clinical years with limited patient contact. Students start their clinical rotations in 3rd-year of MBBS. Therefore, we included only 3rd-year students, both males and females, in this study. Non-probability convenience sampling technique was employed for data collection.

Questionnaire: A self-administered questionnaire was distributed among 3rd-year medical students as developed by Thomas Jefferson University after permission was taken from the Center for Research in Medical Education and Health Care Thomas Jefferson University Sidney Kimmel Medical College. Students were asked to specify their age, gender, and future specialty preference. Specialty preferences were categorized into people-oriented, technology-oriented, and undecided as given in table 1^{10,11}. Students then had to fill the 20-items Jefferson Scale of Empathy Medical Student version (JSE-S).

Measurement of empathy: JSE-S was employed to assess the empathy of medical students in our study. It is a validated tool to calculate the empathy of medical students¹² and is used across the globe. It has been translated into many languages^{10,13,14} including Urdu¹⁵ which is the national language of Pakistan. The original version of the scale in English language was utilized

because all participants understood the English. Besides JSE-S, Jefferson scale of empathy has 2 other versions namely, Health Professionals version (HP-version) for practicing physicians, and Health Profession Students' version (HPS-version) for paramedical students and other health profession students except those of medicine. S-version contains 20 items each of which is scored on a 7-point Likert scale. Of them, 10 items are positively scored with 1 being strongly disagree and 7 being strongly agree. The other 10 items are scored in opposite order with 1 being strongly agree and 7 being strongly disagree. All the items' scores are summed up to obtain a total empathy score which has the range of 120 i.e., 20 score as possible minimum and 140 score as possible maximum. As the mean score increases so does the empathy. The items are further categorized into three domains i.e., perspective taking (10 directly scored items), compassionate care (8 reverse scored items), and standing in the patient's shoes (2 reverse scored items). If a returned questionnaire is missing answers for more than 4 items it is excluded. While missing answers for 4 or fewer items are considered and their values calculated from the average of the filled items scores.

Data analysis: Data were analyzed using the software S.P.S.S version 24.0 for Windows (SPSS Inc, Chicago, IL, USA). Quantitative results were displayed as mean \pm standard deviation while qualitative as frequencies and percentages. Descriptive statistics were calculated for the JSE-S score. Cronbach's alpha was additionally determined to observe the internal consistency of the 20 items. The value greater than 0.9 was interpreted as excellent, greater than 0.8 as good, and greater than 0.7 as acceptable¹⁶. Comparison of JSE-S scores between groups of age and gender was done using independent sample t-test while among specialty preference groups was drawn using one-way analysis of variance (ANOVA) with post hoc test using the Turkey HSD test. A p value of < 0.05 was taken as statistically significant. T-tests and ANOVA were also done for sub-scales of JSE-S i.e., perspective taking, compassionate care, and standing in patient shoes for age, gender, and specialty preference. Cohn's d and eta squared (η^2) were calculated as effect sizes to inspect the practicability of those findings which were statistically significant. For a d of 0.2 effect size was considered as small, for 0.5 as medium, and for 0.8 as large effect size. However, eta squared (η^2) value was taken as small effect size if $\eta^2 = 0.01$, medium effect size if $\eta^2 = 0.06$, and large effect size if $\eta^2 = 0.14$ ¹⁷.

RESULTS

A total of 152 survey forms were distributed among the students. One hundred twenty-nine forms were returned (84.9% response rate). After the invalid form's exclusion, 123 were finally analyzed. The demographics of participants are

presented in Table 1.

Descriptive statistics of JSE-S, including score intervals with frequencies, central tendency and dispersion (mean, SD, range, and percentiles), and distribution (skewness and kurtosis) are displayed in Table 2. A Cronbach's coefficient of 0.82 for the 20-items scale was suggestive of good internal consistency of the scale for our sample. Overall, the mean empathy score recorded was 103.24 ± 15.83 . Among the items, the highest mean recorded was of item no. 20, i.e., 6.16 ± 1.37 , which is regarding their belief about the importance of empathy in medical treatment. On the other hand, the lowest mean was of item no. 18, i.e., 2.85 ± 1.73 , which is about how much a physician should allow himself to be influenced by his personal bond with the patient or their family members.

Across the age groups, participants younger than 22 years had a significantly higher mean empathy than equal to or older than 22 years ($p = 0.004$, two-tailed). Their mean difference was 9.18 (95% CI: 3.00 to 15.37) and had an effect size of medium estimate ($d = 0.56$) (Table 3).

The gender difference on JSE-S mean scores showed females had a significantly higher empathy than males ($p < 0.001$, two-tailed) and had an effect size of large estimate ($d = 0.99$) with a mean difference of -13.91 (95% CI: -18.89 to -8.93) (table 4).

Variables	n (%)
Age	
< 22 years	90 (73.2)
≥ 22 years	33 (26.8)
Gender	
Male	65 (52.8)
Female	58 (47.2)
Future Specialty Preference	
^a People-oriented specialties	57 (46.3)
^b Technology-oriented specialties	48 (39.0)
Undecided	18 (14.6)
^a People-oriented specialties: Internal medicine and medicine sub-specialties including; cardiology, hematology/oncology, general internal medicine, gastroenterology, endocrinology, nephrology, rheumatology, & obstetrics/gynecology, emergency medicine, neurology, general pediatrics, psychiatry, sports medicine, dermatology. ^b Technology-oriented specialties: Surgery & surgical sub-specialties including; cardiothoracic, general surgery, transplant surgery, vascular surgery, & orthopedic surgery, ophthalmology, ENT, plastic surgery, neurosurgery. (10,11)	

Table 1: Demographics of the survey participants

As given in Table 4, the analysis of variance (ANOVA) showed no significant difference in JSE-S mean scores among the three specialty preference groups ($p = 0.08$). However, when ANOVA was done separately for genders, the scores varied significantly among the specialty preference groups for females ($p = 0.02$, η^2

Score interval	Frequency (%)	Cumulative frequency (%)
≤ 60	2 (1.6)	2 (1.6)
61-70	2 (1.6)	4 (3.3)
71-80	7 (5.7)	11 (8.9)
81-90	17 (13.8)	28 (22.8)
91-100	20 (16.3)	48 (39.0)
101-110	26 (21.1)	74 (60.2)
111-120	32 (26.0)	106 (86.2)
121-130	16 (13.0)	122 (99.2)
131-140	1 (0.8)	123 (100.0)
Statistics	value	
Mean \pm SD	103.24 \pm 15.83	
Possible range	120 (20-140)	
Actual range	79 (53-132)	
Skewness	-0.631	
SE skewness	0.218	
Kurtosis	0.306	
SE kurtosis	0.433	
25 th percentile	93	
50 th percentile	104	
75 th percentile	115	
Cronbach's α	0.82	
N	123	

Table 2: Score distribution, descriptive statistics, central tendency and dispersions for JSE-S

	Age		Age	
	< 22 years	≥ 22 years	Male	Female
n	90	33	65	58
Mean \pm SD	105.7 \pm 14.2	96.52 \pm 18.13	96.7 \pm 16.8	110.6 \pm 10.6
df	121		109.49	
t	2.94		-5.53	
p-value (2-tailed)	0.004		< 0.001	
Cohen's d	0.56		0.99	

Table 3: Comparison of empathy across age and gender

= 0.12) while for males it was non-significant ($p = 0.38$). Further, when post-hoc comparisons were done, the underlying significant difference for females was between technology-oriented specialties and undecided groups ($p = 0.02$) with a mean difference of 13.28 (95% CI: 1.92 to 24.65). The difference between people-oriented specialties and undecided was, however, marginally significant ($p = 0.07$) with a mean difference of 10.27 (95% CI: 0.59 to 21.13).

The mean and Standard deviation of the 3 domains of JSE-S and their comparisons across age, gender, and specialty preference are given in Table 5. Among the age and gender groups, younger than 22 years and females had significantly higher perspective taking (age: $p = 0.01$; gender: $p < 0.001$) and compassionate care

(age: $p = 0.004$; gender: $p < 0.001$). For specialty preference groups, no significant difference was seen for these 2 domains ($p = 0.053$, $p = 0.28$, respectively). Differences in mean values of standing in the patient's shoes were non-significant across age ($p = 0.67$), gender ($p = 0.48$), and specialty preference ($p = 0.88$).

DISCUSSION

This study assessed the empathy level of 3rd-year medical students with the help of JSE-S version. Although we did not include other years to see the transition in empathy during medical school. Instead, we only focused on 3rd-year students as studies have shown them the lowest empathic⁷. Observed Cronbach's α of 0.82 supports the reliability of the scale for our

sample. The 20-item questionnaire has good internal consistency as reported in previous studies 0.88^{10} , 0.83^{18} , 0.80^{19} .

In previous studies across Pakistan, 3rd-year empathy as recorded with JSE-S is found to be 87.71 ± 13.94 (Lahore)⁷, 101.49 ± 12.90 (Sukkur)³, 86.15 ± 16.7 (Karachi)²⁰. These values are comparatively lower than our obtained empathy (103.24 ± 15.83). However, when compared with the empathy of foreign medical students i.e., 109.1 ± 11.8 (United States)⁹, 109.75 ± 11.39 (Bangladesh)¹⁰, 104.5 ± 13.9 (Malaysia)²¹, our participants had a lower empathy than them. These differences could possibly be attributed to variation in sample selection, educational systems, curriculum demands, working environment, cultural factors and spiritual beliefs.

	Total			Males			Females		
	Group 1	Group 2	Group 3	Group 1	Group 2	Group 3	Group 1	Group 2	Group 3
n	57	48	18	25	28	12	32	20	6
Mean \pm SD	105.37 \pm 15.32	99.27 \pm 16.09	107.06 \pm 15.29	98.72 \pm 17.86	93.36 \pm 16.97	100.17 \pm 14.09	110.56 \pm 10.66	107.55 \pm 10.38	120.83 \pm 4.07
Between groups df	2			2			2		
Within groups df	120			62			55		
F-value	2.62			0.98			3.96		
p-value	0.08			0.38			0.02		
Group 1 = People-oriented specialties Group 2 = Technology-oriented specialties Group 3 = Undecided									

Table 4: Summary results of one-way ANOVA of empathy scores across specialty-preference groups

	Perspective taking Mean \pm SD	Compassionate care Mean \pm SD	Standing in the patient's shoes Mean \pm SD
Total	54.38 \pm 10.12	40.46 \pm 7.24	8.40 \pm 2.80
Age			
< 22 years	55.78 \pm 9.17	41.59 \pm 6.70	8.33 \pm 2.68
\geq 22 years	50.58 \pm 11.64	37.36 \pm 7.84	8.58 \pm 3.15
p-value	0.01	0.004	0.67
Cohen's d	0.50	0.58	0.08
Gender			
Male	50.69 \pm 11.17	37.75 \pm 7.47	8.23 \pm 3.07
Female	58.52 \pm 6.79	43.48 \pm 5.64	8.59 \pm 2.48
p-value	< 0.001	< 0.001	0.48
Cohen's d	0.85	0.86	0.13
Specialty preference			
People-oriented specialties	55.79 \pm 8.91	41.09 \pm 7.69	8.49 \pm 3.01
Technology-oriented specialties	51.69 \pm 11.76	39.19 \pm 6.39	8.40 \pm 2.56
Undecided	57.11 \pm 7.44	41.83 \pm 7.76	8.11 \pm 2.87
p-value	0.053	0.28	0.88
Eta squared	0.05	0.02	0.002

Table 5: Domains of JSE-S and their comparisons across age, gender, and specialty preferences

The age difference in the empathy score has been found multiple times to be insignificant as shown by Aye SZ et al.²¹ and others. Narang R et al. has obtained similar results even in dental students²⁴. In contrast to these studies, our findings showed that empathy was significantly decreasing for the older age group compared to younger group. Although, we did not find any study that look into the details of the age vs. empathy association, one reason for the decline with age can be a result of going into a higher class⁶ but since the participants were only from 3rd-year the confounding effect of the changing class has been avoided.

In the literature available, the majority studies have found females to have more empathic nature than males^{3,7,9}. There are only a few studies which show otherwise such as by Hussain AA²⁰ and other²³. Our results also agreed with the majority of studies since females scored greater than males. The reason females have higher empathizing ability is explained by both biological factors such as their genetic architecture and social factors such as expected gender roles which make them more cognizable of emotions²⁵. Moreover, females consider interpersonal relations more important than males²⁶.

Although some studies have found no significant effect of specialty preference over empathy^{10,21}, those students who express interest in specialties requiring continuous encounter with patients, classified as people-oriented specialties usually are inclined to have higher empathy towards their patients as compared to technology-oriented and other specialties^{9,18}. In contrast to these studies, our results showed highest empathy among those students who had not yet decided their preference for any future specialty i.e., undecided specialty > people-oriented specialties > technology-oriented specialties. This provides very important data that could be utilized by medical educationists and trainers to have a targeted approach during empathy learning as part of the medical curriculum. However, it is also important to note that this difference was still insignificant in our findings.

While comparing 3 domains of JSE-S across gender groups, females had significantly higher perspective taking ($p < 0.001$) and compassionate care ($p < 0.001$) than males but the difference was non-significant for Standing in the patient's shoes ($p = 0.48$). The study of Hussain AA, found higher perspective taking for females but males dominated the compassionate care and Standing in the patient's shoes as compared to our study²⁰. In addition, we did not see significant variation in the 3 domains across the specialty preference groups ($p = 0.053$, $p = 0.28$, and $p = 0.88$, respectively). In contrast, Hussain AA found the undecided group to have higher scores in all the³ domains of JSE-S than the other specialty preference groups.

CONCLUSION

We found that overall empathy of 3rd-year students in our study is

comparatively greater than other parts of the country. There are variations in empathy across age and gender groups. Specialty preference was found to have little effect on empathy. However, the association of empathy with specialty preference among females is interesting. Integration of empathy into the learning objectives of medical education is necessary because of its clear importance for the doctor and patient⁸. Targeted programs focusing on interpersonal skills, analyzing audio-, video-tape encounters with patients, studying literature and arts, and importance of role models should be taught to them⁹. Also, the variation in empathy among age groups should be looked into because of literature gap.

Acknowledgments: The authors are very grateful to Dr. Farooq Ahmed, Director Medical Education KMC for his general support through each and every step of writing this article

Conflict of Interest: The authors declare no conflict of interest.

Funding: It was a non-funded research work

REFERENCES

1. Stepien KA, Baernstein A. Educating for empathy. *Journal of general internal medicine*. 2006 May;21(5):524-30.
2. Kerasidou A, Horn R. Making space for empathy: supporting doctors in the emotional labour of clinical care. *BMC medical ethics*. 2016 Dec;17(1):1-5.
3. Mirani SH, Shaikh NA, Tahir A. Assessment of clinical empathy among medical students using the Jefferson Scale of Empathy-Student Version. *Cureus*. 2019 Feb 28;11(2).
4. Javaeed A, Rasheed AA, Manzoor A, Ain QT, D COSTA PR, Ghauri SK. Empathy scores amongst undergraduate medical students and its correlation to their academic performance. *Journal of Advances in Medical Education & Professionalism*. 2022 Apr;10(2):99.
5. Shi M, Du T. Associations of emotional intelligence and gratitude with empathy in medical students. *BMC medical education*. 2020 Dec;20(1):1-8.
6. Hizomi Arani R, Naji Z, Moradi A, Shariat SV, Mirzamohamadi S, Salamati P. Comparison of empathy with patients between first-year and last-year medical students of Tehran University of Medical Sciences. *BMC medical education*. 2021 Dec;21(1):1-7.
7. Shaheen A, Mahmood MA, Zia-ul-Miraj M, Ahmad M. Empathy levels among undergraduate medical students in Pakistan, a cross sectional study using Jefferson scale of physician empathy. *J Pak Med Assoc*. 2020 Jul 1;70(7):1149-53.
8. Riess H, Kelley JM, Bailey RW, Dunn EJ, Phillips M. Empathy training for resident physicians: a randomized controlled trial of a neuroscience-informed curriculum. *Journal of general internal medicine*. 2012 Oct;27(10):1280-6.
9. Hojat M, Vergare MJ, Maxwell K, Brainard G, Herrine SK, Isenberg GA, Veloski J, Gonnella JS. The devil is in the third

year: a longitudinal study of erosion of empathy in medical school. *Academic medicine*. 2009 Sep 1;84(9):1182-91.

10. Mostafa A, Hoque R, Mostafa M, Rana M, Mostafa F. Empathy in undergraduate medical students of Bangladesh: psychometric analysis and differences by gender, academic year, and specialty preferences. *International Scholarly Research Notices*. 2014;2014.
11. Chen D, Lew R, Hershman W, Orlander J. A cross-sectional measurement of medical student empathy. *Journal of general internal medicine*. 2007 Oct; 22:1434-8.
12. Hojat M, Mangione S, Nasca TJ, Cohen MJ, Gonnella JS, Erdmann JB, Veloski J, Magee M. The Jefferson Scale of Physician Empathy: development and preliminary psychometric data. *Educational and psychological measurement*. 2001 Apr;61(2):349-65.
13. Paro HB, Daud-Gallotti RM, Tibério IC, Pinto R, Martins MA. Brazilian version of the Jefferson Scale of Empathy: psychometric properties and factor analysis. *BMC Medical Education*. 2012 Dec;12(1):1-7.
14. Bilgel N, Ozcakir A. Turkish version of the Jefferson Scale of Empathy psychometric properties. *European Scientific Journal*. 2017;13(20):101-11.
15. Muhammad SR. Establishing the construct validity and Internal consistency for Urdu version of Jefferson Scale of Empathy-S (JSE-S). medRxiv. 2020 Jun 9:2020-06.
16. Gliem JA, Gliem RR. Calculating, interpreting, and reporting Cronbach's alpha reliability coefficient for Likert-type scales. Midwest research-to-Practice Conference in Adult, Continuing, and community education.
17. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Frontiers in psychology*. 2013 Nov 26; 4:863.
18. Hojat M, DeSantis J, Shannon SC, Mortensen LH, Speicher MR, Bragan L, LaNoue M, Calabrese LH. The Jefferson Scale of Empathy: a nationwide study of measurement properties, underlying components, latent variable structure, and national norms in medical students. *Advances in Health Sciences Education*. 2018 Dec; 23:899-920.
19. Hojat M, Gonnella JS. Eleven years of data on the Jefferson scale of empathy-medical student version (JSE-S): proxy norm data and tentative cutoff scores. *Medical Principles and Practice*. 2015;24(4):344-50.
20. Hussain AA. Empathy among Students of a Public-Sector Medical University: A Cross-Sectional Study. *Journal of Clinical Epidemiology and Toxicology*. SRC/JCET-109. 2020:3.
21. Aye SZ, Sein HH, Nyunt MK, Min Y. Comparative study of Malaysian Medical Students' Empathy Scores Between Preclinical and Clinical Training. *Education in Medicine Journal*. 2022.
22. Javed MQ. The evaluation of empathy level of undergraduate dental students in Pakistan: a cross-sectional study. *Journal of Ayub Medical College Abbottabad*. 2019 Jul 10;31(3):402-6.
23. Iqbal S, Bilal K, Ahmad W, Nazir U, Javed Z, Khan S, Zafar S, Mukhtar F. Empathy among medical students: A cross-sectional survey. *Journal of Ayub Medical College, Abbottabad: JAMC*. 2020 Oct 1;32(4):S681-5.
24. Narang R, Mittal L, Saha S, Aggarwal VP, Sood P, Mehra S. Empathy among dental students: a systematic review of literature. *Journal of Indian Society of Pedodontics and Preventive Dentistry*. 2019 Oct 1;37(4):316-26.
25. Greenberg DM, Warriar V, Abu-Akel A, Allison C, Gajos KZ, Reinecke K, Rentfrow PJ, Radecki MA, Baron-Cohen S. Sex and age differences in "theory of mind" across 57 countries using the English version of the "Reading the Mind in the Eyes" Test. *Proceedings of the National Academy of Sciences*. 2023 Jan 3;120(1):e2022385119.
26. Ardenghi S, Rampoldi G, Bani M, Strepparava MG. Attachment styles as predictors of self-reported empathy in medical students during pre-clinical years. *Patient education and counseling*. 2020 May 1;103(5):965-70

CONFLICT OF INTEREST

Author declared no conflict of interest

GRANT SUPPORT & FINANCIAL DISCLOSURE

Author declared no specific grant for this research from any funding agency in the public, commercial or non-profit sectors

AUTHORS CONTRIBUTIONS

MIM&MK: Conception, Design of the work, Data collection, and Drafting, Reviewed, Final approval, Agreement to be accountable.

HA&NU: Conception, Design of the work, Acquisition, Data Analysis, and Drafting, Reviewed, Final approval, Agreement to be accountable.



This is an Open Access article distributed under the terms of the Attribution Non-Commercial 4.0 International